

Finding outputs of a two-step function with decimals that models a real-world situation: Two variable equation: Worksheet 8.3

Name Date Score

1. John rented a truck for one day. There was a base fee of \$40. And there was an additional charge of \$4 for each mile driven. The total cost in dollars for driving x miles is given by the following function $C(x) = 40 + 4x$. What is the total cost if John drove 25 miles?
2. Oceania Bike Rentals charges 20 dollars plus 4 dollars an hour for renting a bike. Total amount to be paid is given by the function $T(h) = 20 + 4h$, where h is the number of hours. What total amount Sandy has to pay if he rented a bike for 6 hours?
3. The sum of three consecutive numbers is given by the function $3n + 3$ where n is the smallest number. If the smallest of the three numbers is 59 what is the sum of the three consecutive numbers?
4. A small pond has 800 liters of water to start with. Water is added to the pond at the rate of 50 liters per minute. If water is added for t minutes the total capacity of the pond T is given by $T = 800 + 50t$. Find the capacity of the pond after a period of 25 minutes.
5. The money Nancy had spent on books is given by the function $T = 20 + 9b$ where b is the cost of one book. How much did she spend if cost of each book is \$6?
6. The number of books Ron has is given by the function $T(x) = 0.5x + 25$, where x is the number of books he has initially. How many books does he have now, if $x = 20$?
7. The amount that Jessica has is given by the function $A(x) = 0.5x + 15$, where x is her allowance in dollars. What is the amount she has if her allowance is \$24?
8. The number of students who went on a trip to the zoo is given by the function $N(s) = 8s + 21$ where s is the number of students traveling in each bus. How many students went to the zoo if $s = 25$?



Solutions: Worksheet 8.3

- The number of cards Tom has is given by the function $T(k) = 0.75k + 14$, where k is the number cards in a pack. Find the number of cards Tom has if $k = 48$?
- The sum of three consecutive odd numbers is given by the function $S(n) = 3n + 6$, where n is the smallest of the three given odd numbers. If $n = 27$, what is the sum of the three consecutive odd numbers?

Solutions: Worksheet 8.3

- \$140
- \$44
- 180
- 2050 liters
- \$74
- 35 books
- \$27
- 221 students
- 50 cards
- 87

